

NSA-226-62  
Code-1

N 63 17215

UNCOLLECTED PRELIMINARY DATA

SYNTHESIS OF PORPHINE-LIKE SUBSTANCES  
DURING CHEMICAL EVOLUTION

OTS PRICE

XEROX

\$

MICROFILM

\$

ANTON SZUTKA

DEPARTMENT OF CHEMISTRY

UNIVERSITY OF DETROIT

4001 W. McNichols Road

Detroit 21, Michigan

## INTRODUCTION

It is already a well established theory that chemical evolution, i.e. evolution on the molecular level, has preceded the appearance of the first living cell on the earth. Current concepts suggest that a relatively simple primitive atmosphere, consisting of methane, ammonia, water, and also probably some molecular hydrogen under influence of ultraviolet radiation, electrical discharges and high energy radiations has been transformed into more complicated compounds (1, 2, 3). It has been generally accepted that the temperature at the time of this transformation was less than 100°C. The raw material thus produced, accumulated in pools of water that condensed, forming an organic millieu, i.e. the nonliving mixture of carbon compounds which is generally considered as a prerequisite of the origin of life. Some estimates made by Urey (2) indicate that the primitive oceans might have contained as much as 10 per cent of dissolved organic compounds. The synthesis of these raw materials continued until a state of equilibrium was reached, in which the action of available energy sources produced as much of the raw material as it destroyed. In this period of time only relatively simple molecules could be formed because the more complicated and labile ones would be destroyed by the energetic ultraviolet radiation reaching the surface of the earth, unscreened by the ozone layer, which now protects the earth.

The above theories of Dparin (1) and Uray (2) have been subjected to rigid laboratory tests by Miller (4,5), Pavlovskaya and Pasynskii (6), Abelson (7), Oro (8), Hasselstrom (9), Paschke (10), and others (11, 12). This laboratory work has produced various products of biochemical importance, such as amino acids, aliphatic acids, polyhydroxy compounds, aldehydes and urea. Subjecting these simple molecules to further action of ultraviolet radiation or mild heating, produced a variety of complex organic molecules. Such higher order reactions, for example, result in production of polypeptides from amino acids, as demonstrated by Fox (13, 14), Akabori (15), Doty (16), and others (17), in actual laboratory experiments.

In the reviewed literature, most of the authors (1, 18, 19, 20) dealing with the problem of the origin of life, are convinced that at some stage of chemical evolution, porphyrins must have developed. However, until recently there was no laboratory evidence that porphyrins or porphine-like substances could be synthesized from precursors which were available during the period of chemical evolution on earth.

It has been shown in our previous paper (21), that porphine-like substances, specifically  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  -tetraphenylporphines, can be synthesized from simple precursors using  $\gamma$ -radiation for activation purposes.

This paper presents another observation having direct bearing on the formation of the organic millieu, namely the synthesis of porphine-like substances from simple precursors.

## EXPERIMENTAL PROCEDURE

A system, consisting of 3 ml of freshly distilled pyrrole, 6 ml of benzaldehyde, and 4 ml of water, was placed in a 250 ml beaker, and then irradiated, at a distance of 25 cm with ultra-violet light, using a 100 Watt Hanovia Utility Model Lamp. The radiation transmitted through the filter was 45 per cent at 2500 Å, 90 per cent at 3000 Å and the output of the lamp was  $1.5 \times 10^{15}$  quanta per sec per  $\text{cm}^2$  or  $5 \times 10^{16}$  quanta per sec per area of the mixture exposed toward radiation. The doses were measured by chemical actinometry, using an uranyl oxalate actinometer according to Forbes and Heidt (22). After exposure, the reddish-brown solution was diluted with approximately 20 ml of chloroform and washed with 20 ml of distilled water. After separation in a separatory funnel, the aqueous layer was discarded, and the chloroform layer was transferred into a beaker charged with approximately 20 g of anhydrous sodium sulfate to remove the excess of water. The sodium sulfate was then filtered off and discarded, while the filtrate was made up to 50 ml volume with chloroform and stored for the quantitative determination of porphines.

The final product separation was accomplished by column chromatography, using activated alumina (F-20) and Florosil (60-100 mesh) as adsorbents and fresh chloroform as eluent. Two chromatographic treatments on alumina and three to four chromatographic separations on Florosil yielded pure product. The porphines appeared in

eluent and collections of eluent were continued until no Soret band appeared in the effluent. The final spectrum of the product was recorded on a Bausch and Lomb Spectronic 505 recording spectrophotometer, using benzene as a solvent.

A typical spectrum of compound isolated from irradiated mixture and the reference spectrum of the compound synthesized by the method of Ball, et al. (23), are presented in Fig. 1. Both spectra are identical with those obtained by Thomas et al. (24), and Dorrough et al. (25). For quantitative estimation of the yield, the molar extinction coefficient of  $18.7 \times 10^3$  at 515 mμ maximum was used, as suggested by Thomas and Martell (24). Identity of the compound was further established by the formation of a zinc chelate, according to the procedure of Rothmund and Menotti (26), and Dorrough et al. (25). Both spectra, the reference zinc chelate and the chelate of the compound isolated from the irradiated mixture are identical, as shown in Fig. 2.

The same mixture of pyrrole, benzaldehyde and water when placed in the dark without irradiation, also produced porphines. The isolation and quantitative determination of porphines was done by the same methods used for irradiated mixtures.

If, instead of water, pyridine was used as a solvent in the system, and the mixture was then irradiated for two hours, no porphine-like substances could be detected by previously used methods. Aging of the mixture for ten days produced only traces of porphines.

Fig. 3 represents the total yields of porphines isolated from the stored mixtures, plotted against time of storage. Each line in this graph represents different irradiation periods.

## DISCUSSION

Literature dealing with the origin of life, particularly the role of porphines and porphyrins in life's origin have raised several important questions. As yet, they are unanswered; the future can and must answer them.

In our present study we have shown that porphine-like structures can be synthesized in the presence of oxygen from precursors which were available in the very earliest stage of chemical evolution. Aldehydes were found in products of the action of electrical discharges on the Urey atmosphere in Miller's experiment (4, 5). Pyrroles and pyrrolidines were easily formed from ammonia, acetylene, and other unsaturated hydrocarbons by simple catalysis or under influence of ultraviolet radiation (1). In addition, Lichtin (27) found pyrroles as the major product of the reaction of active nitrogen with 1,3-butadienes.

The results of this study also indicate that the presence of water, or rather the suspension of the organic matter in water, increases the yield of porphine like substances considerably. Oparin (1), and Haldane (3), have already indicated the possibility

of evolution of living matter from the pools rich in organic matter, rather than on solid surfaces. The concentration of organic matter could be quite high, if part of the water solvent evaporated, as the case might be in lakes and lagoons.

The increase in the yield of porphine-like substances, on standing, shows that a process of autocatalysis takes place. This process, postulated by Calvin (19), now can be supported by experimental evidence.

There is a general agreement among many authors that synthesis of porphyrins is a necessary step for the origin of life. However, the question of the time of the appearance of porphyrins is subject to discussion. Gaffron (18), and Calvin (19), support the idea that porphyrins were already active in the very earliest quasi-living organic structures. Miller and Urey (5), consider this not a necessity and suggest that porphyrins may have arisen during the evolution of primitive organisms. Strughold and Ritter (28), consider pre-existing stores of oxygen, produced by photochemical dissociation of water, as a prerequisite for the formation and development of chlorophyll, while the heme types of porphyrins were formed about 1 to 1.5 billion years later.

At the present time, it seems to be impossible to determine at what stage of chemical evolution, porphine-like substances developed. In the present living processes, porphine-like substances find a great variety of uses, such as fermentation, respiration, and photosynthesis. The metabolism of sulfates, iron, nitrogen, hydrogen,

and oxygen also require porphyrins. In addition, an interesting observation was made by Baltrop (19c) on the formation of pyrophosphates from phosphates in presence of ferrous protoporphyrins. All these facts would suggest that porphyrins evolved prebiologically in order to facilitate the evolution of seemingly living organic structures up to the present state. This would place the appearance of porphyrins in the oxygen free period of chemical evolution.

On the other hand, after the stores of simpler organic compounds reached the saturation point, in the first stage of chemical evolution, there was need for a reduction of the energy level of the source in order to continue chemical evolution to a more complicated nature. Photolysis of water and the formation of an ozone layer cut out ultra-violet radiation completely, while visible light itself did not provide enough energy for chemical synthesis. With the aid of porphyrins and especially their metal chelates, chemical transformations such as hydrogen transfer or oxidation could occur.

Simultaneously with this event, hydrogen peroxide appeared. If it remained in contact with organic substances, the results would be fatal. Therefore, in order to continue chemical evolution, it was necessary that a mechanism be developed for the destruction of vast amounts of hydrogen peroxide. Calvin (19d), has already pointed out that incorporation of ferric iron into a heme-type molecule would increase the catalytic activity of iron by a factor of  $10^3$  and the addition of certain protein arrangements by an additional factor of  $10^5$ .

The last two arguments indicate that with the transition of



the reductive atmosphere into oxidative, an immediate need was created for the porphine-like substances. This need served as an evolutionary selection pressure for the synthesis of porphine-like substances.

This work is supported in part by the Grant N S G - 226-62 from the National Aeronautics and Space Administration


## REFERENCES

1. (a) A.I. Oparin, The Origin of Life on the Earth (translated by A. Synge), 3rd ed., Academic Press, New York, 1957; (b) Life, its Nature, Origin and Development (translated by A. Synge), Academic Press, New York, 1962.
2. H.C. Urey, The Planets, Yale University Press, New Haven, Connecticut, 1952.
3. J.B.S. Haldane, Collection of Essays on the Origin of Life, Penguin Books, Ltd., London, 1954.
4. (a) S.L. Miller, A production of amino acids under possible primitive earth conditions. *Science* 117, 528-529 (1953); (b) Production of some organic compounds under possible primitive earth conditions. *J. Am. Chem. Soc.* 77, 2351-2361 (1955); (c) The formation of organic compounds on the primitive earth. *Ann. N.Y. Acad. Sci.* 69, 260-274 (1957); (d) The mechanism of synthesis of amino acids by electric discharges. *Biochim. et Biophys. Acta* 23, 480-489 (1957).
5. S.L. Miller and H.C. Urey, Organic Compound Synthesis on the Primitive earth. *Science* 130, 245-251 (1959).
6. T.E. Pavlovskaya and A.G. Pasyanskii, The original formation of amino acids under action of ultraviolet rays and electric discharges. In Proceedings of the First International Symposium on the Origin of Life on the Earth. A.I. Oparin, et al. ed. pp. 151-157, Pergamon Press, London, (1959).

7. P.H. Abelson, Amino acids formed in "primitive atmosphere".  
Science 124, 935 (1956)
8. (a) J. Oro, Mechanism of synthesis of adenine from hydrogen cyanide under possible primitive earth conditions. Nature 191, 1193-1194 (1961); (b) J. Oro and C.L. Guidry, Direct Synthesis of polypeptides. I. Polycondensation of glycine in aqueous ammonia. Arch. Biochem. Biophys. 93, 166-171 (1961); (c) J. Oro and S.S. Kamat, Amino acid synthesis from hydrogen cyanide under possible primitive earth conditions. Nature 190, 442-443 (1961); (d) J. Oro and A.P. Kimball, Synthesis of purines under possible primitive earth conditions. I. Adenine from hydrogen cyanide. Arch. Biochem. Biophys., 94, 217-227 (1961).
9. (a) T. Hasselstrom and M.C. Henry, New Synthesis of Oxalic Acid. Science, 123, 1038-1039 (1956); (b) T. Hasselstrom, M.C. Henry, and B. Murr, Synthesis of Amino Acids by Beta Radiation. Science, 125, 350-351 (1957).
10. R. Paschke, R.W.H. Chang, and O. Young, Probable role of gamma irradiation on origin of life. Science 125, 881 (1957).
11. (a) K. Bahadur, Photosynthesis of amino acids from paraformaldehyde and potassium nitrate. Nature 173, 1141 (1954); (b) K. Bahadur and R.R. Srivastava, Effect of catalysts and the period of irradiation on the photosynthesis of amino acids in the mixture of para-formaldehyde and potassium nitrate. Zh. Obshchei Khimii 31, 3017-3020 (1961).
12. A. Palm and M. Calvin, Primordial Organic Chemistry. I. Compounds resulting from electron irradiation of  $C^{14}H_4$ . J. Am. Chem. Soc., 84, 2115 (1962).

13. (a) S.W. Fox, Evolution of protein molecules and thermal synthesis of biochemical substances. *Am. Scientist* 44, 347-362 (1956); (b) S.W. Fox, How did life begin? *Science* 132, 200-208 (1960).
14. S.W. Fox and K. Harada, Thermal copolymerization of amino acids in the presence of phosphoric acids. *Arch. Biochem. Biophys.* 66, 281-285 (1960).
15. S. Akabori, On the origin of the fore-protein. In *Proc. of the First International Symposium on the Origin of Life on the Earth* (A.I. Oparin et al., eds. pp. 189-196, Pergamon Press, London, (1959).
16. P. Doty, K. Imahori, and E. Klemperer, The solution, properties and configurations of a polyampholytic polypeptide, copoly-L - lysine - L - glutamic acid. *Proc. Nat. Acad. Sci.* 44, 424-431 (1958).
17. (a) G. Ferrari and R. Cultrera, Sul meccanismo della sintesi fotochimica di amino acidi. Nota I. *Gazzetta Chim. Ital.* 90, 1522-1529 (1960); (b) G. Ferrari and R. Cultrera, Sul meccanismo della sintesi fotochimica di amino acidi. Nota II. *Gazz. Chim. Ital.* 90, 1637-1644 (1960)
18. H. Gaffron, The origin of life. *Perspectives in Biol. Med.* 3, 163-212 (1960)
19. (a) M. Calvin, Evolution of enzymes and the photosynthetic apparatus. *Science* 130, 1170-1174 (1959); (b) M. Calvin, The Chemistry of life. *Chem. Eng. News* 39, 96-104 (May 22, 1961); (c) M. Calvin, The origin of life on earth and elsewhere. *Perspectives in Biol. Med.* 5, 399-422 (1962); (d) M. Calvin, Evolution of photosynthetic mechanisms, *Perspectives in Biol. Med.* 5, 147-172 (1962).

# SPECTRA IN BENZENE SOLUTION



## Zn - Chelate SPECTRA in BENZENE

